

# NeoMed<sup>®</sup> 325

## (neomycin sulfate)

### Soluble Powder

#### Antibacterial

For Use in Cattle, Swine, Sheep, Goats & Turkey

ANADA 200-050, Approved by FDA



MADE IN USA

### DESCRIPTION

NeoMed<sup>®</sup> 325 is an odorless powder containing 71.4 g neomycin sulfate equivalent to 50 g neomycin per 100 g of product.

### INDICATIONS

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in cattle (excluding veal calves), swine, sheep and goats. For the control of mortality associated with *Escherichia coli* organisms susceptible to neomycin sulfate in growing turkeys.

### BENEFITS

- **Safeguard:** Approved for treatment of growing turkeys with no withdrawal period
- **Versatile:** Provides treatment for a wide variety of species
- **Soluble:** Dissolves readily in drinking water
- **Convenient:** Available in 100 g and 200 g packets plus space saving resealable 50 lb. bulk pails
- **Safe:** Approved by FDA



### PACKAGING

LIST NO.	UNIT PACKAGE	CASE SIZE
1NE0013	100 g (3.5 oz) pouch	50
1NE0010	200 g (7 oz) pouch	25
1NE0007	50 lb (22.7 kg) pail	1

### WARNING

DISCONTINUE TREATMENT PRIOR TO SLAUGHTER BY AT LEAST THE NUMBER OF DAYS LISTED BELOW FOR APPROPRIATE SPECIES:

Cattle .....	1 day
Swine and Goats .....	3 days
Sheep .....	2 days
Turkeys .....	0 days

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

A milk discard period has not been established for this product in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older.

#### Bimeda, Inc.

One Tower Lane, Suite 2250

Oakbrook Terrace, IL 60181 USA

Toll Free Tel. 888-524-6332 • Toll Free Fax. 877-888-7035

Email: sales@bimedaus.com

See reverse side for Administration and Dosage.



# NeoMed® 325

(neomycin sulfate)

**Antibacterial**

**For Use in Cattle, Swine, Sheep, Goats & Turkey**

**Soluble Powder**

ANADA 200-050, Approved by FDA

**NOT FOR HUMAN USE. FOR ANIMAL USE ONLY.**

**KEEP THIS AND ALL DRUGS OUT OF REACH OF CHILDREN**

**Restricted Drug (California) – Use Only as Directed**

**DESCRIPTION:**

Add to drinking water - Not for use in liquid supplements.

**INDICATIONS:**

For the treatment and control of colibacillosis (bacterial enteritis) caused by *Escherichia coli* susceptible to neomycin sulfate in cattle (excluding veal calves), swine, sheep and goats. For the control of mortality associated with *Escherichia coli* organisms susceptible to neomycin sulfate in growing turkeys.

**DOSAGE AND ADMINISTRATION:**

Administer to cattle (excluding veal calves), swine, sheep and goats at a dose of 10 mg neomycin sulfate per pound of body weight per day in divided doses for a maximum of 14 days. Administer to turkeys at a dose of 10 mg neomycin sulfate per pound of body weight per day for 5 days.

**HERD/FLOCK TREATMENT:**

100 g (3.5 oz) NeoMed® 325 will treat 7150 pounds of body weight. Estimate the total number of pounds of body weight of the animals to be treated and administer 3.5 oz for each 7150 pounds. The product should be added to the amount of drinking water estimated to be consumed in 12-24 hours. Provide medicated water as the sole source of water each day until consumed, followed by non-medicated water as required. Fresh medicated water should be prepared each day.

**INDIVIDUAL ANIMAL TREATMENT:**

To provide 10 mg neomycin sulfate per pound of body weight, mix one (1) level teaspoon in water or milk for each 160 pounds body weight. Administer daily either as a drench in divided doses or in the drinking water to be consumed in 12-24 hours.

**DRINKING WATER:**

**Swine** - Use the amount of NeoMed® 325 indicated below in 256 gallons of water, or in two gallons of stock solution used in proportioner set to meter one ounce per gallon.

- Pigs weighing 25 to 50 Pounds .....7 oz
- Pigs weighing 50 to 100 Pounds .....10.5 oz
- Pigs weighing over 100 Pounds.....14 oz

Manufactured by:  
Bimeda, Inc.  
Le Sueur, MN 56058  
www.bimeda.com

NeoMed® is a registered trademark of Bimeda, Inc.  
629-13 NeoMed TS 08/2013

**CAUTION:**

To administer the stated dosage, the concentration of neomycin required in medicated water must be adjusted to compensate for variation in age and weight of animal, the nature and severity of disease signs, and environmental temperature and humidity, each of which affects water consumption.

If symptoms persist after using this preparation for 2 or 3 days, consult a veterinarian. If symptoms such as fever, depression, or going off feed develop, oral neomycin is not indicated as the sole treatment since systemic levels of neomycin are not obtained due to low absorption from the gastrointestinal tract.

**IMPORTANT:**

Treatment should continue 24 to 48 hours beyond remission of disease symptoms, but not to exceed a total of 14 consecutive days for cattle, swine, sheep and goats and 5 days for turkeys. Animals not drinking or eating should be treated individually by drench.

**WARNING:**

**DISCONTINUE TREATMENT PRIOR TO SLAUGHTER BY AT LEAST THE NUMBER OF DAYS LISTED BELOW FOR APPROPRIATE SPECIES:**

- Cattle .....1 day
- Swine and Goats.....3 days
- Sheep.....2 days
- Turkeys .....0 days

A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

A milk discard period has not been established for this product in lactating dairy cattle. Do not use in female dairy cattle 20 months of age or older.

Securely reseal bulk pails and tightly fold packets to keep out moisture.

**STORAGE:** Store at controlled room temperature 20° C - 25° C (68° F - 77° F), with excursions permitted to 15° C - 30° C (59° F - 89° F).

**HOW SUPPLIED:**

NeoMed® 325 is supplied in 100 g (3.5 oz) pouch, packed 50 per pail; 200 g (7.0 oz) pouch, packed 25 per pail; and 50 lb. resealable bulk pail.

To obtain an MSDS or for assistance, contact Bimeda, Inc. at 1-888-524-6332.

TAKE TIME



OBSERVE LABEL DIRECTIONS

[www.BimedaUS.com](http://www.BimedaUS.com)